arXiv:1001.3467v1 [hep-ph] 20 Jan 2010

Two pion production in electron-polarized proton scattering

A.I. Ahmadov*

JINR-BLTP, 141980 Dubna, Moscow region, Russian Federation and

Institute of Physics, Azerbaijan National Academy of Sciences, Baku, Azerbaijan

E.A. Kuraev† and Yu.M. Bystritskiy‡

JINR-BLTP, 141980 Dubna, Moscow region, Russian Federation

Process of two pion production in electron - polarized proton scattering is investigated. In Weizsäcker-Williams approximation the differential spectral distributions and the spin-momentum correlations are considered. The spin correlation effects caused by ρ-meson widths are estimated to be of order of several percents. Both channels of $\pi^+\pi^-$ and $\pi^+\pi^0$ creation are considered. The effects of intermediate excited baryons are not considered. The spectral distributions on pion energy fractions in polarized and unpolarized cases are presented analytically and numerically.

I. INTRODUCTION

We consider the kinematical region of small angles electron scattering, which corresponds to the fragmentation region of the initial polarized proton. Besides we are interesting the case of small invariant masses of the final system consisting of the nucleon and the pair of mesons supposed to be not exceeding the mass of baryon resonances. In this case the intermediate state with ρ-meson becomes important. Apart of Weitzsaker-Williams (WW) enhancement [1–3], which is associated with "large logarithmic factor" ($\ln\left(s^2/\left(M_p^2 m_e^2\right)\right) \sim 20$), an additional enhancement associated with Breit-Wigner character of the intermediate ρ-meson propagator takes place when considering the total cross section as well as differential distributions. The rather large width of ρ-meson induce the measurable one-spin asymmetries [4, 5] then initial proton is polarized transversally to the electron beam direction (laboratory

---

*Electronic address: ahmadov@theor.jinr.ru

†Electronic address: kuraev@theor.jinr.ru

‡Electronic address: bystr@theor.jinr.ru

frame is implied).

Our paper is organized as follows. We start the general consideration in Section II where we consider the processes

$$\begin{aligned} e^-(p_1)+p(p) &\to e^-(p_1')+p(p')+\pi^+(q_+)+\pi^-(q_-); \\ e^-(p_1)+p(p) &\to e^-(p_1')+n(p')+\pi^+(q_+)+\pi^0(q_0). \end{aligned}$$

The definite channels of $\pi^+\pi^-$ production and $\pi^+\pi^0$ production are considered in Sections III, IV correspondingly. In Section V we gather the final results and demonstrate some numerical estimations for experimental observables.

Comparison with data obtained at HERMES facility are given.

II. GENERAL CONSIDERATION

The elastic scattering kinematics is implied

$$p_1^2=p_1'^2=m_e^2, \qquad p^2=p'^2=M^2; \qquad q_\pm^2=q_0^2=m^2. \tag{1}$$

Besides we consider the case of small angles of electron scattering:

$$-q^2=-(p_1-p_1')^2 \ll s, \qquad s=2pp_1 \gg M^2. \tag{2}$$

So for HERMES facility we have $s \approx 30-40$ GeV^2. Matrix element in Born approximation has a form

$$M^{ep\to en2\pi}=\frac{4\pi\alpha G}{q^2}\left[\bar{u}(p_1')\gamma_\mu u(p_1)\right]J_\nu(q)g^{\mu\nu}, \tag{3}$$

where $J_\nu(q)$ is the hadron current

$$J_\nu(q)=\bar{u}(p')O_\nu u(p), \tag{4}$$

which obeys the current conservation condition $J_\lambda(q)q^\lambda=0$, operator O_ν will be defined in Sections III, IV. The quantity G is a product of ρ-meson coupling constants $G=g_{\rho\pi\pi}g_{\rho NN}$. For the peripheral kinematics the Sudakov parametrization of 4-vectors of the problem is useful [6–8]. For this aim we introduce two almost light-like 4 vectors $\tilde{p}_1$ and $\tilde{p}$:

$$\begin{aligned} &\tilde{p}_1=p_1-p\frac{m_e^2}{s}; \qquad \tilde{p}=p-p_1\frac{M^2}{s}, \\ &\tilde{p}_1^2=O\left(m_e^4\frac{M^2}{s^2}\right); \qquad \tilde{p}^2=O\left(m_e^4\frac{M^4}{s^2}\right), \\ &2\tilde{p}\tilde{p}_1=s; \qquad 2p\tilde{p}=M^2; \qquad 2p_1\tilde{p}_1=m_e^2. \end{aligned} \tag{5}$$

And the corresponding parametrization reads as

$$\begin{aligned} q &= \alpha\tilde{p}+\beta\tilde{p}_1+q_\perp; \\ q_\pm &= x_\pm\tilde{p}+\beta_\pm\tilde{p}_1+q_{\pm\perp}; \\ q_0 &= x_0\tilde{p}+\beta_0\tilde{p}_1+q_{0\perp}; \\ p' &= x\tilde{p}+\beta\tilde{p}_1+q_\perp, \end{aligned} \tag{6}$$

where the orthogonality conditions ($p_1 a_\perp = p a_\perp = 0$) takes place. The 4-vectors $a_\perp$ have only spatial components, besides it's z-axis component is zero (we choice as a z-axis the direction of motion of initial electron $a_\perp = (0,0,\vec{a})$, where $\vec{a}$ is a 2-dimensional vector transversal to the z-axis. Thus we have

$$a_\perp^2 = -\vec{a}^2 < 0. \tag{7}$$

The conservation law and final particles mass-shell conditions give the following relations:

$$\begin{aligned} s\beta_\pm &= \frac{1}{x_\pm}\left(\vec{q}_\pm^{\,2}+m^2\right), \\ s\beta' &= \frac{1}{x}\left(\vec{p}^{\,\prime 2}+M^2\right); \\ \vec{q} &= \vec{p}^{\,\prime}+\vec{q}_+ +\vec{q}_-, \\ x+x_-+x_+ &= 1, \\ \beta+\frac{M^2}{s} &= \beta_+ +\beta_- +\beta', \end{aligned} \tag{8}$$

for the subprocess of $\pi^+\pi^-$ photoproduction: $\gamma^*(q)+p(p)\to p(p')+\pi^+ +\pi^-$. The similar relations for the subprocess $\gamma^*(q)+p(p)\to n(p')+\pi^+ +\pi^0$ have a form

$$\begin{aligned} s\beta_+ &= \frac{1}{x_+}(\vec{q}_+^{\,2}+m^2); \\ s\beta_0 &= \frac{1}{x_0}(\vec{q}_0^{\,2}+m^2); \\ s\beta' &= \frac{1}{x}(\vec{p}^{\,\prime 2}+M^2); \\ \vec{q} &= \vec{p}^{\,\prime}+\vec{q}_+ +\vec{q}_0, \\ x+x_+ +x_0 &= 1, \\ \beta+\frac{M^2}{s} &= \beta_+ +\beta_0 +\beta'. \end{aligned} \tag{9}$$

Significant simplification comes from the Gribov's form of the metric tensor $g^{\mu\nu}$ in photon propagator [6]:

$$g^{\mu\nu} = g_\perp^{\mu\nu} + \frac{2}{s}\left[\tilde{p}^\mu\tilde{p}_1^\nu + \tilde{p}^\nu\tilde{p}_1^\mu\right] \approx \frac{2}{s}\tilde{p}^\mu\tilde{p}_1{}^\nu. \tag{10}$$

In the kinematics we are considering the only $(2/s)\tilde{p}^\mu \tilde{p}_1^\nu$ component of this tensor is relevant. Other terms give the contributions which are suppressed by factors

$$O\left(\frac{M^2}{s}\right) \ll 1. \tag{11}$$

We also systematically omit the terms of order

$$O\left(\frac{m^2}{M^2}\right) \sim 0.02. \tag{12}$$

The terms of this order of magnitude (which are systematically omitted below) determine the accuracy of our consideration.

Thus matrix element has a form:

$$\begin{aligned} M^{ep\to eN2\pi} &= \frac{8\pi s\alpha}{q^2} N_e M_h^i, \qquad i = \pi^+\pi^-, \pi^+\pi^0, \\ M_h^i &= \frac{1}{s} p_1^\nu J_\nu(q), \\ N_e &= \frac{1}{s}\left[\bar{u}(p_1')\gamma_\mu u(p_1)\right]\tilde{p}^\mu. \end{aligned} \tag{13}$$

It is easy to see that

$$\sum |N_e|^2 = 2, \tag{14}$$

and that the quantity N_h does not depend on s in the large s limit.

Let us now parameterize the phase volume of the final particles:

$$d\Gamma = (2\pi)^{-8} \frac{d^3 p_1'}{2E_1'} \frac{d^3 p'}{2E'} \frac{d^3 q_1}{2E_1} \frac{d^3 q_2}{2E_2} \delta^4(p + p_1 - p' - p_1' - q_1 - q_2). \tag{15}$$

We use below the on-mass shell relation $(i = +, -, 0)$

$$\frac{d^3 q_i}{2E_i} = d^4 q_i' \delta\left(q_i^2 - m\right) = \frac{s}{2} dx_i d\alpha_i d^2\vec{q}_i \delta\left(s x_i \alpha_i \vec{q}_i^{\,2} - m^2\right) = \frac{1}{2x_i} dx_i d^2\vec{q}_i. \tag{16}$$

Introducing the additional unit factor

$$1 = d^4 q \; \delta^4\left(p_1 - p_1' - q\right) = \frac{s}{2} d\alpha \; d\beta \; d^2\vec{q} \; \delta^4\left(p_1 - p_1' - q\right) \tag{17}$$

we put the phase volume to the form

$$d\Gamma = (2\pi)^{-8} \frac{dx_1 dx_2}{8 s x x_1 x_2} d^2\vec{q} d^2\vec{q}_1^{\,2} d^2\vec{q}_2. \tag{18}$$

The channel of $\pi^+\pi^-$ production corresponds to $\vec{q_1} \equiv \vec{q_-}$, $x_1 \equiv x_-$, $\vec{q_2} \equiv \vec{q_+}$, $x_2 \equiv x_+$; while the channel of $\pi^+\pi^0$ production corresponds to $\vec{q_1} \equiv \vec{q_0}$, $x_1 \equiv x_0$, $\vec{q_2} \equiv \vec{q_+}$, $x_2 \equiv x_+$. In phase volume transformation we used the mass shell conditions for scattered electron

$$p_1'^2 - m_e^2 = (p_1 - q)^2 - m_e^2 = s\alpha\beta - \vec{q}^2 - s\alpha - m_e^2\beta = 0, \tag{19}$$

and the conservation laws (8) and (9). Specific attention should be payed to expressing the transfer momentum squared q^2. Using the scattered electron mass shell condition (19) we can write it as

$$q^2 = s\alpha\beta - \vec{q}^2 = -\frac{1}{1-\beta}\left(\vec{q}^2 + m_e^2\beta^2\right) \approx -\left(\vec{q}^2 + m_e^2\beta^2\right). \tag{20}$$

The quantity $s\beta$ can be expressed in terms of invariant mass squared of final hadronic system:

$$s_0 = s\beta = (p' + q_1 + q_2)^2 + \vec{q}^2 - M^2. \tag{21}$$

This quantity in Weitzsaker-Williams approximation can be written as

$$\begin{aligned}(s_0)_{\vec{q}=0} &= \frac{M^2}{xx_1x_2}\left[x_1x_2(1-x) + \nu^2 x(1-x)^2 + x_1(1-x_1)\chi_2 + \right.\\ &+ \left. x_2(1-x_2)\chi_1 + 2x_1x_2\chi\right] \equiv \frac{s_1M^2}{xx_1x_2},\end{aligned} \tag{22}$$

where

$$\nu = \frac{m}{M}; \qquad \chi_i = \frac{\vec{q}_i^2}{M^2}, \qquad \chi = \frac{\vec{q_1}\vec{q_2}}{M^2}.$$

We see that the transferred momentum q is the space-like 4-vector with non-zero module. One can see that the module of the matrix element tends to zero in the limit then transferred momentum q goes to zero. To be convinced let us write the quantity N_h as

$$M_h = \frac{1}{s}p_1^\nu J_\nu(q) = \frac{1}{s_0}(q - q_\perp)^\nu J_\nu(q) = -\frac{1}{s_0}q_\perp^\nu J_\nu(q), \tag{23}$$

where we have used the current conservation condition.

Here and further we restrict ourselves only by the approximation of a "large logarithm" (i.e. so called Weitzsaker-Williams approximation). For this aim we perform the integration over the transferred momentum using the relation

$$\int \frac{d^2\vec{q}\ \vec{q}_i\vec{q}_j}{\pi(\vec{q}^2 + m_e^2\beta^2)^2} = \frac{1}{2}\delta_{ij}(L-1),$$

where L is the "large logarithm":

$$L = \ln \frac{Q^2 s^2}{m_e^2 s_0^2} \approx \ln \frac{s^2}{m_e^2 M^2} \approx 23. \tag{24}$$

Writing the matrix element of the pions photoproduction sub-process $\gamma^*(q) + p \to N\pi_1\pi_2$ as

$$M_h^i = \frac{1}{s_0} \bar{u}(p') \hat{O}_i u(p), \qquad i = \pi^+\pi^-, \pi^+\pi^0, \tag{25}$$

and using the density matrix of a polarized proton as $u(p,a)\bar{u}(p,a) = (\hat{p} + M)(1 - \gamma_5 \hat{a})$, where 4-vector a is the polarization vector of the initial proton (i.e. $p \cdot a = 0$), we obtain for the differential cross section:

$$\frac{\pi^2 M^2 d\sigma_i}{dx_1 dx_2 d^2\vec{q}_1 d^2\vec{q}_2} = \frac{\alpha^2 G^2}{64 M^2 \pi^3}(L-1) x x_1 x_2 \frac{S_i}{s_1^2}, \tag{26}$$

where

$$S_i = \frac{1}{4M^2} Tr \left[\left(\hat{p}' + M \right) \hat{O}_i \left(\hat{p} + M \right) \left(1 - \gamma_5 \hat{a} \right) \hat{\bar{O}}_i \right]. \tag{27}$$

We note that for HERMES conditions the factor $\sigma_0 = \alpha^2 G^2 (L-1)/(64 M^2 \pi^3)$ have a rather large value $\sigma_0 \approx 2 \cdot 10^{-30}$ cm^2, i.e. of order of two microbarns. The explicit expressions for the quantity S_i for both processes considered here are given in Appendices A, B.

III. MATRIX ELEMENT OF $\pi^+\pi^-$ PRODUCTION ON A PROTON

The matrix element of $\pi_+\pi_-$ mesons production can be presented as a sum of two separately gauge-invariant contributions

$$M_{\pi_+\pi_-} = M_1 + M_2, \tag{28}$$

First term describes the subprocess of creation of pion pair by (virtual) photon and (virtual) ρ-meson

$$\begin{aligned} M_1 \; &= \; \frac{G}{q_1^2 - M_\rho^2} \left[\bar{u}(p') \gamma^\rho u(p) \right] \times \\ &\times \left\{ \frac{\left(\left(2q_- - q \right), e \right) \left(-2q_+ + q_1 \right)_\rho}{(q - q_-)^2 - m^2} + \frac{\left(2q_- - q_1 \right)_\rho \left(\left(-2q_+ + q \right), e \right)}{(q - q_+)^2 - m^2} - 2e_\rho \right\}, \end{aligned} \tag{29}$$

where $e = e(q)$ is the polarization vector of virtual photon, $q_1 = q_+ + q_- - q$, $q_2 = q_+ + q_-$. Second term describes the subprocess of Compton scattering on a proton with emission of ρ-meson which subsequently decays into pion pair

$$M_2 = \frac{G}{q_2^2 - M_\rho^2} \left[\bar{u}(p') \left\{ \hat{e} \frac{\hat{p}' - \hat{q} + M}{(p'-q)^2 - M^2} \hat{v} + \hat{v} \frac{\hat{p} + \hat{q} + M}{(p+q)^2 - M^2} \hat{e} \right\} u(p) \right], \tag{30}$$

where $v = -q_+ + q_-$. Here we omit the QED coupling α since it is already taken into account in (26).

Both contributions ($M_1$and M_2) satisfy the gauge condition: they turn to zero when use the mass shell conditions and replace $e(q) \to q$.

The matrix element (28) can be written in the form

$$M_{\pi_+\pi_-} = \frac{sG}{s_0} \left[\bar{u}(p') \hat{O}_{+-} u(p) \right], \tag{31}$$

where we left only quadratic over $\vec{q}$ terms and operator $\hat{O}_{+-}$ has a form:

$$\hat{O}_{+-} = A\hat{q}_+ + B\hat{q}_- + C\hat{q}_\perp + D\hat{v} + E\hat{q}_\perp \hat{p}_1 \hat{v} + F\hat{v}\hat{p}_1\hat{q}_\perp, \tag{32}$$

where

$$A = \frac{4x^2 x_+}{s_1 D_\rho} \frac{\vec{q}\vec{q}_-}{M^2}; \qquad B = \frac{4x^2 x_-}{s_1 D_\rho} \frac{\vec{q}\vec{q}_+}{M^2}; \qquad D = \frac{2x_-^2 x_+^2}{s_1 D_\rho} \frac{\vec{p}'\vec{q}_-}{M^2};$$

$$C = \frac{2x}{D_\rho}; \qquad E = \frac{x_+ x_-}{s x D_\pi}; \qquad F = \frac{x_+ x_-}{s D_\pi},$$

and the ρ-meson and pion propagators denominators are

$$\begin{aligned} D_\rho &= x\mu^2 + \nu^2(1-x)^2 + \chi_1 + \chi_2 + 2\chi, \\ D_\pi &= -\mu^2(1 - i\gamma_\rho)x_+x_- + x_1^2\chi_2 + x_2^2\chi_1 - 2x_+x_-\chi + \nu^2(1-x)^2, \end{aligned}$$

$$\gamma_\rho = \frac{\Gamma_\rho}{M_\rho}, \qquad \mu = \frac{M_\rho}{M}, \qquad \nu = \frac{m}{M}.$$

IV. MATRIX ELEMENT OF $\pi^+\pi^0$ PRODUCTION ON A PROTON

For the subprocess $\gamma^*(q) + p(p) \to n(p')\pi_+(q_+)\pi_0(q_0)$ the amplitude M_{+0} contains besides the Compton-like amplitude

$$M_1 = \frac{G}{(q_+ + q_0)^2 - M_\rho^2} \left[\bar{u}(p')(\hat{q}_0 - \hat{q}_+) \frac{\hat{p} + \hat{q} + M}{(p+q)^2 - M^2} \hat{e} u(p) \right], \tag{33}$$

also three terms corresponding to conversion of virtual photon and virtual ρ-meson to pion pair:

$$\begin{aligned}
M_2 &= G\frac{\bar{u}(p')\gamma_\rho u(p)}{\left((q_+ + q_0)^2 - M_\rho^2\right)\left((q_+ + q_0 - q)^2 - M_\rho^2\right)} \\
&\times \left[ev_+\left(-q_+ - q_0 - q\right)_\rho + v_{+\rho}\left(2q_+ + 2q_0 - q\right)e + e_\rho v_+\left(2q - q_+ - q_0\right)\right], \\
M_3 &= -G\frac{\bar{u}(p')(-\hat{q}_+ + \hat{q}_0 + \hat{q})u(p)e(-2q_+ + q)}{\left((q_+ + q_0)^2 - M_\rho^2\right)\left((q_+ - q)^2 - m^2\right)}; \\
M_4 &= G\frac{\bar{u}(p')\hat{e}u(p)}{(q_+ + q_0 - q)^2 - M_\rho^2}, \qquad v = v_+ = -q_+ + q_0.
\end{aligned}$$

It can be checked that the sum

$$M_{+0} = M_1 + M_2 + M_3 + M_4, \tag{34}$$

obeys the gauge condition. Writing this amplitude in the form similar to (31) leads us to

$$M_{+0} = \frac{sG}{s_0}\left[\bar{u}(p')\hat{O}_{+0}u(p)\right], \tag{35}$$

where:

$$\begin{aligned}
\hat{O}_{+0} &= -\frac{\hat{v}_+\hat{p}_1\hat{q}_\perp}{sD_\pi} + \frac{1}{D_\rho D_\pi}\left[-2\beta\hat{p}_1\vec{q}\vec{v}_+ + 2\hat{v}_+\vec{q}(\vec{q}_+ + \vec{q}_0) + s_1(x_+ - x_0)\hat{q}_\perp\right] - \\
&- \frac{1}{s_1 D_\rho x_+}\left[s_1 x_+\hat{q}_\perp + 4\vec{q}\vec{q}_+\hat{q}_0\right],
\end{aligned} \tag{36}$$

where the expressions for s_1, D_ρ, D_π can be obtained from the relevant expressions for process of $\pi_+\pi_-$ production (see Section III) by replacement $\vec{q}_-, x_- \to \vec{q}_0, x_0$.

V. RESULTS

The differential cross section of pion pair electroproduction processes which we are considering here can be obtain using Equ. (26). In case of $\pi^+\pi^-$ pair production we have to insert the quantity S_{+-} into this formula (26). This quantity has a form:

$$\begin{aligned}
S_{+-} &= (1 - P_\pm)\left\{\frac{-2\frac{1}{M}\left[\vec{q_+}\vec{a}\right]_z \cdot ImD_\pi}{D_\rho|D_\pi|^2 s_1}A + \frac{4ReD_\pi M^2}{s_1^2 D_\rho|D_\pi|^2}D\right\} + \\
&+ (1 + P_\pm)\left\{\frac{2M^2}{D_\rho^2 s_1^2}C + \frac{2M^2}{s_1^2|D_\pi|^2}F\right\},
\end{aligned} \tag{37}$$

x_+ / x_-	0.2	0.3	0.4	0.5	0.6	0.7
0.2	0.268	0.359	0.362	0.312	0.250	0.138
0.3	0.337	0.613	0.708	0.620	0.410	
0.4	0.351	0.712	0.899	0.808		
0.5	0.315	0.625	0.792			
0.6	0.240	0.406				
0.7	0.137					

Tab. I: The function $F_{+-}^{\mathrm{unp}}\left(x_+, x_-\right)$ (defined in (41)) is presented for different values of final pions energy fractions x_+ and x_- for $\pi^+\pi^-$ production case.

where permutation operator $P_\pm$ acts on any function $F\left(x_+, x_-, \vec{q}_+, \vec{q}_-\right)$ in the following manner

$$P_\pm F\left(x_+, x_-, \vec{q}_+, \vec{q}_-\right) = F\left(x_-, x_+, \vec{q}_-, \vec{q}_+\right) \tag{38}$$

and the quantities A, C, D, F in (37) are given in Appendix A.

In case of $\pi^+\pi^0$ pair production we have to insert the quantity S_{+0} into this formula (26). This quantity has a form:

$$\begin{aligned} S_{+0} &= -\frac{2ImD_\pi}{s_1 D_\rho^2 |D_\pi|^2}\left[\frac{\left[\vec{q}_+, \vec{a}\right]_z}{M}\left(A_1 + A_2 D_\rho\right) + \frac{\left[\vec{q}_0, \vec{a}\right]_z}{M} B\right] + \frac{2}{s_1 D_\rho^2} C + \\ &+ \frac{2}{D_\rho |D_\pi|^2} D + \frac{4ReD_\pi}{s_1 D_\rho^2 |D_\pi|^2} E + \frac{2}{D_\rho |D_\pi|^2} F + \frac{4ReD_\pi}{D_\rho |D_\pi|^2} G + \frac{2}{|D_\pi|^2} H, \end{aligned} \tag{39}$$

where quantities A_1, A_2, B, C, D, E, F, G, H are given in Appendix B.

Dalitz-distribution (the distribution over final pions energy fractions) can be written in the following form

$$\frac{d\sigma}{dx_1 dx_2} = \frac{\alpha^2 G^2}{64 M^2 \pi^3}\left(L - 1\right) F_i^{\mathrm{unp}}\left(x_1, x_2\right), \tag{40}$$

where function $F_i^{\mathrm{unp}}\left(x_1, x_2\right)$

$$F_i^{\mathrm{unp}}\left(x_1, x_2\right) = \int \frac{d^2\vec{q}_1}{\pi}\frac{d^2\vec{q}_2}{\pi} x x_1 x_2 \frac{S_i}{s_1^2}. \tag{41}$$

This function in case of $\pi^+\pi^-$ production is presented in Table I, while in case of $\pi^+\pi^0$ production is in Table II.

x_+ / x_0	0.2	0.3	0.4	0.5	0.6	0.7
0.2	56.325	23.089	2.897	0.418	0.052	0.003
0.3	23.166	14.824	5.237	0.421	0.016	
0.4	2.876	5.063	0.990	0.071		
0.5	0.415	0.419	0.067			
0.6	0.051	0.016				
0.7	0.003					

Tab. II: The function $F^{\mathrm{unp}}_{+0}(x_+, x_0)$ (defined in (41)) is presented for different values of final pions energy fractions x_+ and x_0 for $\pi^+\pi^0$ production case.

In the experimental setup when we fix the azimuthal angle ψ between the proton polarization vector $\vec{a}$ and transverse momentum of one of pions $\vec{q_+}$, i.e. $\psi = (\vec{a}, \vec{q_+})$, some polarization dependent contributions appear:

$$\int d\left|\vec{q_1}\right| \int \frac{d^2 q_2}{\pi} x x_1 x_2 \frac{S_i}{s_1^2} = |\vec{a}| \sin\psi \; F_i^{\mathrm{pol}}(x_1, x_2) + F_i^{\mathrm{unp}}(x_1, x_2). \tag{42}$$

And thus in experiment the asymmetry [4, 5]

$$A(x_1, x_2) = \frac{F_i^{\mathrm{pol}}(x_1, x_2)}{F_i^{\mathrm{unp}}(x_1, x_2)}, \tag{43}$$

can be measured. This asymmetries for both channels are presented in Figs. 1, 2 as a function x_+ for fixed values of x_- or x_0.

We should note that the meaning of quantities x, x_1, x_2 is different in laboratory reference frame and in the center of mass of the initial particles (c.m. frame). In c.m. frame they are the energy fractions of the jet consisting of the recoil proton and pions which obey the conservation laws

$$x + x_1 + x_2 = 1, \qquad x_i = \frac{2E_i}{\sqrt{s}}, \qquad \frac{2m}{\sqrt{s}} < x_i < 1, \qquad \frac{2M}{\sqrt{s}} < x < 1. \tag{44}$$

In Laboratory frame, keeping in mind the explicit form of the light-like vectors $\tilde{p}$ and $\tilde{p}_1 \approx p_1$:

$$\tilde{p} = \frac{M}{2}(1, -1, 0, 0), \qquad p_1 = E(1, 1, 0, 0),$$

we have for energies:

$$E_i = \frac{M}{2} x_i + \frac{\vec{q}_i^{\,2} + m^2}{2 M x_i}, \qquad E' = \frac{M}{2} x + \frac{(\vec{q_1} + \vec{q_2})^2 + M^2}{2 M x},$$



Fig. 1: The asymmetry $A\left(x_{+}, x_{-}\right)$ (defined in (43)) as a function of x_{+} with x_{-} fixed to definite values.



Fig. 2: The asymmetry $A\left(x_{+}, x_{0}\right)$ (defined in (43)) as a function of x_{+} with x_{0} fixed to definite values.

$$M + \frac{s_1}{2Mxx_1x_2} = E_1 + E_2 + E', \qquad \frac{m}{M} < x_i < 1.$$

The values of 3-momenta squares are

$$\begin{aligned} \mathbf{q}_i^2 &= \vec{q}_i^{\,2} + \left[-\frac{M}{2}x_i + \frac{\vec{q}_i^{\,2} + m^2}{2Mx_i}\right]^2, \\ \mathbf{p}^2 &= (\vec{q}_1 + \vec{q}_2)^2 + \left(-\frac{M}{2}x + \frac{(\vec{q}_1 + \vec{q}_2)^2 + M^2}{2Mx}\right)^2. \end{aligned}$$

We specifically emphasize that in left part of these equalities are 3-dimensional vectors and in the right part – the two dimensional vectors (their components perpendicular to the initial electron direction are zero).

Dalitz-plot distributions for unpolarized case (see Fig. 1 and 2, while $\sigma_0 \approx 2\mu b$) seems to be accessible in experiments of HERMES. The one-spin asymmetries (see (43)) have a value of several percents and as well can be measured. Preliminary comparison of these quantities are in qualitative agreement with data obtained at HERMES [9].

Appendix A: The production of $\pi^+\pi^-$ pair

Here we present the the explicit expressions for the values A, C, D, F for process $\pi_+\pi_-$ production (see (37)):

$$A = A_1\chi_+ + \frac{1}{2}A_3\chi_- + \frac{1}{2}A_4 s_1 + \frac{1}{2}A_5; \tag{A1}$$

$$\begin{aligned} C = {} & C_1\chi_+^2\chi_- + \frac{1}{2}C_3\chi_+\chi_-\chi + C_4\chi_+^2 + \frac{1}{2}C_6\chi^2 + \frac{1}{2}(C_7 + s_1C_8)\chi_+\chi_- + \\ & (C_9 + s_1C_{10})\chi_+\chi + (C_{13}s_1 + C_{14} + C_{15}s_1^2)\chi_+ + \frac{1}{2}(C_{19} + C_{20}s_1^2)\chi + \frac{1}{2}C_{21}s_1^2 + \frac{1}{2}C_{22}s_1^3; \end{aligned} \tag{A2}$$

$$\begin{aligned} D = {} & D_1\chi_+^2\chi_- + D_3\chi_+\chi_-^2 + D_5\chi_+^2\chi + \frac{1}{2}D_7\chi_+\chi_-\chi + D_8\chi_+^2 + \frac{1}{2}\chi^2(D_{10} + s_1D_{11}) + \\ & \frac{1}{2}\chi_+\chi_-D_{12} + (D_{13} + s_1D_{14})\chi_+\chi + (D_{17}s_1 + D_{18} + s_1^2D_{19})\chi_+ + \\ & \frac{1}{2}(D_{23}s_1 + D_{24} + D_{25}s_1^2)\chi + \frac{1}{2}D_{26}s_1^2 + \frac{1}{2}D_{27}s_1^3; \end{aligned} \tag{A3}$$

$$\begin{aligned} F = {} & F_1\chi_+^3 + F_3\chi_+^2\chi_- + F_5\chi_+\chi^2 + F_7\chi_+^2\chi + \frac{1}{2}F_9\chi_+\chi_-\chi + \chi_+^2(F_{10}s_1 + F_{11}) + \\ & \frac{1}{2}\chi^2(F_{14}s_1 + F_{15}) + \frac{1}{2}\chi_+\chi_-(F_{16}s_1 + F_{17}) + (F_{18}s_1 + F_{19})\chi_+\chi + \\ & (F_{22}s_1 + F_{23} + s_1^2F_{24})\chi_+ + \frac{1}{2}(F_{28} + s_1F_{29})\chi + \frac{1}{2}s_1^2F_{30} + \frac{1}{2}s_1^3F_{31}. \end{aligned} \tag{A4}$$

The expressions for coefficients are

$$\begin{gathered} A_1 = -x(1-x)(1-x_-)x_-^3;\;\; A_2 = xx_+x_-(1-x)(2x_+ - x_-(1-x_+)); \\ A_3 = 2x(2-x_-)x_+x_-^2(1-x);\;\; A_4 = x_-^2(1-x)(1+x_+-x_-); \\ A_5 = xx_-^3x_+(1-x)(x_+-x_-). \end{gathered} \tag{A5}$$

$$\begin{aligned} C_1 &= -\frac{2}{x_+}x^4(x_-^2 - x_+(1-x_-)); C_3 = 4x^4(2-x); C_4 = -\frac{2}{x_+}x^4x_-^4; \\ C_6 &= 4x^3; C_7 = -2x^4[x_+x_-(1-x) - 2(x_+^2+x_-^2)]; C_8 = \frac{2x^3}{x_+x_-}(1-x)^2; \\ C_9 &= 4x_-^2x^4(1+x_-); C_{10} = -\frac{2}{x_+}x_-x^3; C_{13} = \frac{2}{x_+}x_-^2x^3(1-x); \\ C_{14} &= 2x_-^4x^4; C_{15} = -\frac{x^2}{x_+}; C_{19} = 4x_-^2x_+^2x^4; \\ C_{20} &= \frac{2}{x_+x_-}x^2(1-x); C_{21} = -x^2(1-x); C_{22} = \frac{x}{x_+x_-}. \end{aligned} \tag{A6}$$

$$\begin{gathered} D_1 = 2(xx_-)^2x_+; D_3 = 2(xx_-x_+)^2; D_5 = x^2x_-^2x_+(1-x_-); \\ D_7 = x_+x_-x^2(x_- - x_+)(2-x); D_8 = x^2x_-^4x_+(1+x_-); \\ D_{10} = -2(xx_+x_-)^2(1-x)(x_- - x_+); D_{11} = xx_+x_-(1+x)(x_- - x_+); \\ D_{12} = -3(xx_-x_+)^2(x_- - x_+); \\ D_{13} = x_-^2x^2x_+[x_-^3 + x_-^2(1-2x_+) - x_-x_+(1+x_+) + 2x_+^2]; \\ D_{14} = \frac{1}{2}xx_-[x_-^3 - x_-^2 + x_-x_+(6-x_+) + x_+(3x_+-2)]; D_{17} = -xx_-^3x_+; \\ D_{18} = -x^2x_+^2x_-^4(x_- - x_+); D_{19} = \frac{1}{2}x_-[x_+ - xx_-]; \\ D_{23} = \frac{1}{2}xx_-x_+(1-x)(1-3x)(x_- - x_+); D_{24} = -(xx_-x_+)^2(x_-^2 + x_+^2)(x_- - x_+); \\ D_{25} = \frac{1}{2}x(1-x)(x_- - x_+); D_{26} = -\frac{1}{2}x_-x_+(1-x)(x_- - x_+); \\ D_{27} = \frac{1}{2}(x_- - x_+). \end{gathered} \tag{A7}$$

$$F_1 = \frac{1}{2}x_-^4 x_+^2 (1 - x_-); F_3 = -\frac{1}{2}x_-^3 x_+^2 [x_-^2 - x_-(1 - x_+) + x_+(2 + x_+)];$$
$$F_5 = 2x_-^4 x_+^3; F_7 = x_+^2 x_-^4 (1 + x_+ - x_-); F_9 = -2(x_- x_+)^3;$$
$$F_{10} = -\frac{1}{2}x_-^3 x_+^2 (2 - x_-); F_{11} = -\frac{1}{2}x_-^4 x_+^3 [x_-(3 - x) - 2x_+]; F_{14} = -2(x_- x_+)^3;$$
$$F_{15} = 2(x_- x_+)^4 (1 - x); F_{16} = \frac{1}{2}(x_- x_+)^2 [x_-^2 + x_+^2 + 2(1 - x)];$$
$$F_{17} = -\frac{1}{2}(x_- x_+)^3 [(1 - x)(x_-^2 + x_+^2) + 2(x_- - x_+)^2]; F_{18} = x_-^3 x_+^2 (x_- - x_+);$$
$$F_{19} = -x_-^4 x_+^3 (3 - x)(x_- - x_+); F_{22} = (x_- x_=)^3 (x_- - x_+); F_{23} = \frac{1}{2}(x_- x_+)^4 (x_- - x_+);$$
$$F_{24} = \frac{1}{4x}x_-^2 x_+ [-x_-^2 (1 - x_-) - x_+^2 (3 - x_-) + 2x_+(1 + x_-^2)]; F_{28} = (x_- x_+)^4 (x_- - x_+)^2;$$
$$F_{29} = -\frac{1}{2x}(x_- x_+)^2 [1 + x^2]; F_{30} = -\frac{1}{4x}(x_- x_+)^2 (1 - x)(x_- - x_+)^2;$$
$$F_{31} = \frac{1}{4x}x_- x_+ (x_- - x_+)^2. \tag{A8}$$

Appendix B: The production of $\pi^+\pi^0$ pair

Here we present the the explicit expressions for scalar coefficients in trace for the process $\pi_+\pi_0$ production (see (39)):

$$A_1 = s_1[\chi_+ a_1 + \chi_0 a_2 + \chi a_3] + s_1^2 a_4;$$
$$A_2 = \chi_+ a_5 + \chi_0 a_6 + \chi a_7 + s_1 a_8 + a_9, \tag{B1}$$

with

$$a_1 = -\frac{1}{x_+}x^2 x_0^3; a_2 = x_+ x_0 x^2; a_3 = 4(x_0 x)^2; a_4 = \frac{1}{x_+}x x_0^2;$$
$$a_5 = x^2 x_0^4; a_6 = x_0 x_+ x^2 (x_+ - x_0 + x_+ x_0); a_7 = -2x_+ x_0^3 x^2,$$
$$a_8 = x(x_+ - x_0); a_9 = x_+ x_0^3 x^2 (x_+ - x_0). \tag{B2}$$

$$B = s_1(b_1\chi_+ + b_2\chi_0 + b_3\chi) + b_4 s_1^2, \tag{B3}$$

with

$$b_1 = -(x_0 x)^2; b_2 = (x x_+)^2; b_3 = -4x_+ x_0 x^2; b_4 = -x x_+. \tag{B4}$$

$$C = c_1\chi_+^2\chi_0 + c_2\chi_0^2\chi_+ + c_3\chi_+\chi_0\chi + c_4\chi_+^2 + (c_5 + s_1 c_6)\chi_+\chi_0 + (c_7 + s_1 c_8)\chi_+\chi +$$
$$(c_9 s_1^2 + s_1 c_{10} + c_{11})\chi_+ c_{12} s_1^2 \chi_0 + (s_1 c_{13} + c_{14}) s_1 \chi +$$
$$+ s_1^2 c_{15} + s_1^3 c_{16}, \tag{B5}$$

with

$$\begin{aligned}
c_1 = -\frac{2x_0^2(x_+ + x_0 - 1)^4}{x_+},\ c_2 = -2(x_+ - 1)(x_+ + x_0 - 1)^2,\ c_3 = 4x_0(x_+ + x_0 - 1)^4,\\
c_4 = -\frac{2x_0^4(x_+ + x_{)} - 1)^4}{x_+},\ c_5 = -2(x_+ - 2)x_0^2(x_+ + x_0 - 1)^4,\\
c_6 = -\frac{(x_+ + x_0 - 1)^3(x_+ + 2x_0)}{x_+},\ c_7 = 4x_0^3(x_+ + x_0 - 1)^4,\ c_8 = \frac{x_0(x_+ + x_0 - 1)^3}{x_+},\\
c_9 = -\frac{(x_+ + x_0 - 1)^2}{4x_+},\ c_{10} = -\frac{x_0^2(x_+ + x_0 - 1)^3(x_+ + 2x_0)}{x_+},\ c_{11} = 2x_0^4(x_+ + x_0 - 1)^4,\\
c_{12} = -\frac{(x_+ + x_0 - 1)^2}{4x_0},\ c_{13} = \frac{(x_+ + x_0 - 1)}{x_+},\ c_{14} = x_+x_0(x_+ + x_0 - 1)^3,\\
c_{15} = -\frac{1}{4}(x_+ + x_0 - 1)^2(x_+ + x_0),\ c_{16} = -\frac{x_+ + x_0 - 1}{4x_+x_0}.
\end{aligned} \tag{B6}$$

$$\begin{aligned}
D = d_1\chi_+^3 + d_2\chi_0^3 + d_3\chi_+^2\chi_0 + d_4\chi_0^2\chi_+ + d_5\chi_+^2\chi +\\
d_6\chi_0^2\chi + d_7\chi^2\chi_+ + d_8\chi^2\chi_0 + d_9\chi_+\chi_0\chi +\\
(s_1d_{10} + d_{11})\chi_+^2 + (s_1d_{12} + d_{13})\chi_0^2 + (s_1d_{14} + d_{15})\chi^2 +\\
(s_1d_{16} + d_{17})\chi_+\chi_0 + (s_1d_{18} + d_{19})\chi_+\chi + (s_1d_{20} + d_{21})\chi_0\chi +\\
(s_1^2d_{22} + s_1d_{23} + d_{24})\chi_+ + (s_1^2d_{25} + s_1d_{26} + d_{27})\chi_0 +\\
(s_1^2d_{28} + d_{29})\chi_+s_1^2d_{30} + s_1^3d_{31},
\end{aligned} \tag{B7}$$

with

$$\begin{aligned}
d_1 = -\frac{1}{2}x^2x_0^2(x_0-1),\; d_2 = -\frac{1}{2}x^2x_+^2(x_+-1),\\
d_3 = -\frac{1}{2}x^2x_0(x_0^2+(x_+-1)x_0+x_+(x_++2)),\\
d_4 = -\frac{1}{2}x^2x_+(x_0^2+(x_++2)x_0+x_+(x_+-1)),\\
d_5 = x^2x_0^2(x_+-x_0+1)x_0^2,\; d_6 = -x^2x_+^2(x_+-x_0-1),\\
d_7 = 2x^2x_+x_0^2,\; d_8 = 2x^2x_+^2x_0,\; d_9 = -2x^2x_+x_0, d_{10} = \frac{1}{2}xx_0(x_++2x_0-2),\\
d_{11} = -\frac{1}{2}x^2x_+x_0^2(x_+(x_0-2)+x_0(x_0+2)), d_{12} = \frac{1}{2}xx_+(2x_++x_0-2),\\
d_{13} = -\frac{1}{2}x^2x_+^2x_0(x_+(x_++2)+x_0(x_0-2)), d_{14} = -2xx_+x_0,\; d_{15} = 2x^2x_+^2x_0^2(1-x),\\
d_{16} = -x((x_+-1)x_0-x_+),\\
d_{17} = -\frac{1}{2}x^2x_+x_0(x_0^3+(x_++2)x_0^2+(x_+-4)x_+x_0+x_+^2(x_++2)),\\
d_{18} = -xx_0(x_+-x_0),\; d_{19} = x^2x_0^2x_+(x_+-x_0)(3-x),\\
d_{20} = xx_+(x_+-x_0),\; d_{21} = -x^2x_+^2x_0(x_+-x_0)(3-x),\\
d_{22} = \frac{1}{4}\left(-\frac{x_0^2}{x_+}-3x_++2\right),\; d_{23} = -\frac{1}{2}xx_+x_0(x_+-x_0)(2+x),\\
d_{24} = \frac{1}{2}x^2x_+^2x_0^2(x_+-x_0)^2,\; d_{25} = \frac{1}{4}\left(-\frac{x_+^2}{x_0}-3x_0+2\right),\\
d_{26} = -\frac{1}{2}x_+xx_0(x_+-x_0)(x-1),\; d_{27} = \frac{1}{2}x^2x_+^2x_0^2(x_+-x_0)^2,\\
d_{28} = -x,\; d_{29} = x^2x_+^2(x_+-x_0)^2,\; d_{30} = -\frac{1}{4}(x_+-x_0)(1-x),\\
d_{31} = \frac{(x_+-x_0)^2}{4xx_+x_0}.
\end{aligned} \tag{B8}$$

$$\begin{aligned}
E = e_1\chi_+^2\chi_0 + e_2\chi_0^2\chi_+ + e_3\chi_0^2\chi + e_4\chi^2\chi_0 + e_5\chi_+\chi_0\chi + e_6\chi_+^2\\
(s_1e_7+e_8)\chi^2 + (s_1e_9+e_{10})\chi_0\chi_+ + (s_1e_{11}+e_{12})\chi\chi_+ +\\
(s_1e_{13}+e_{14})\chi\chi_0 + (s_1^2e_{15}+s_1e_{16}+e_{17})\chi_+ + (s_1e_{18}+e_{19})s_1\chi_0 +\\
(s_1^2e_{20}+s_1e_{21}+e_{22})\chi + s_1^2e_{23} + s_1^3e_{24},
\end{aligned} \tag{B9}$$

with

$$\begin{aligned}
e_1 = -x_0 x^3 (x_0+1),\ e_2 = -x_+ x^3 (x_+-1),\ e_3 = -x_+ x^3 (x_+-1),\\
e_4 = -2x^3 x_+ x_0,\ e_5 = -x x_0 (2x_+ - x_0 - 1),\ e_6 = x^3 x_0^3 (x_0+1),\\
e_7 = 2x^2 x_0,\ e_8 = -2x^3 x_+ x_0^3,\ e_9 = -x^2,\ e_{10} = x x_+ x_0 (x_+ + (x_+-2)x_0),\\
e_{11} = \frac{x_0 x^2 (2x_+ - x_0)}{2x_+},\ e_{12} = -x^3 x_0^3 (2x_+ - x_0 - 1),\ e_{13} = -\frac{1}{2} x^2 (3x_+ + 2x_0 - 2),\\
e_{14} = x_+ x_0 x^3 (x_+ + (x_+-2)x_0),\ e_{15} = \frac{x(x_+ - x_0)}{4x_+},\ e_{16} = -\frac{1}{2} x_0^2 x^2 (x_0+2),\\
e_{17} = x_+ x_0^3 x^3 (x_+ - x_0),\ e_{18} = \frac{x(x_+ - x_0)}{4x_0},\ e_{19} = \frac{1}{2} x^2 x_+^2 x_0,\\
e_{20} = -\frac{x(x_+ - x_0)}{2x_+},\ e_{21} = x^2 x_0 (x_+^2 - x_0^2 + x_0),\ e_{22} = x^3 x_+ x_0^3 (x_+ - x_0),\\
e_{23} = \frac{1}{4} x (x_+ - x_0)(1-x),\ e_{24} = \frac{1}{4}\left(\frac{1}{x_+} - \frac{1}{x_0}\right).
\end{aligned} \tag{B10}$$

$$\begin{aligned}
F = f_1\chi_+^2 + f_2\chi_0^2 + f_3\chi^2 + f_4\chi\chi_0 + f_5\chi_+\chi_0 + f_6\chi_0\chi\\
(s_1 f_7 + f_8)\chi_+ + (s_1 f_9 + f_{10})\chi_0 + s_1\chi f_{11} + s_1^2 f_{12},
\end{aligned} \tag{B11}$$

with

$$\begin{aligned}
f_1 = -\frac{1}{2} x_+ x_0^2 x (x_0 - 1),\ f_2 = -\frac{1}{2} x_+^2 x_0 x (x_+ - 1),\ f_3 = 2x_+^2 x_0^2 x,\\
f_4 = -\frac{1}{2} x_+ x_0 x (x_+^2 + x_0^2 + 1 - x),\ f_5 = x_+ x x_0^2 (x_+ - x_0),\ f_6 = -x_+^2 x_0 x (x_+ - x_0),\\
f_7 = \frac{1}{2} x_0 (x_0^2 + (x_+ - 1)x_+),\ f_8 = \frac{1}{2} x_+^2 x_0^2 x (x_+ - x_0),\ f_9 = \frac{1}{2} x_+ (x_+^2 + (x_0 - 1)x_0),\\
f_{10} = -\frac{1}{2} x_+^2 x_0^2 x (x_+ - x_0),\ f_{11} = x_+ x_0 x,\ f_{12} = \frac{1}{4} x_0^2 x.
\end{aligned} \tag{B12}$$

$$\begin{aligned}
G = g_1\chi^2 + g_2\chi_0\chi_+ + g_3\chi\chi_+ + g_4\chi\chi_0 +\\
(s_1 g_5 + g_6)\chi_+ + s_1\chi_0 g_7 + (s_1 g_8 + g_9)\chi + s_1^2 g_{10},
\end{aligned} \tag{B13}$$

with

$$\begin{aligned}
g_1 = -x_+ x_0^2 x^2,\ g_2 = \frac{1}{2} x_+ x_0 x^2,\ g_3 = \frac{1}{2} x_0^3 x^2,\ g_4 = \frac{1}{2}(x_+ - 1) x_+ x_0 x^2,\ g_5 = \frac{1}{4} x_0^2 x,\\
g_6 = \frac{1}{4} x_+ x_0^3 x^2,\ g_7 = -\frac{1}{4} x_+^2 x,\ g_8 = \frac{1}{2}(x_+ - x_0) x_0 x,\ g_9 = -\frac{1}{2} x_+ x_0^3 x^2,\ g_{10} = \frac{1}{4}(x_+ - x_0)
\end{aligned} \tag{B14}$$

$$H = h_1\chi_+ + h_2\chi_0 + h_3\chi + h_4 s_1 + h_5, \tag{B15}$$

with

$$h_1 = -\frac{1}{4}x_+ x_0^2(x_0^2 + x_+(x_0 - 1)),\ h_2 = -\frac{1}{4}x_+^2 x_0(x_+^2 + (x_+ - 1)x_0),\ h_3 = -\frac{1}{2}x_+^2 x_0^2 x,$$
$$h_4 = \frac{x_+(x_+ - x_0)^2 x_0}{4x},\ h_5 = \frac{1}{4}x_+^2 x_0^2(x_+ - x_0)^2. \tag{B16}$$

[1] C. Weizsaecker, Zs. f. Phys. **Bd 88**, 612 (1934).

[2] E. J. Williams, Phys. Rev. **45**, 729 (1934).

[3] A. Akhiezer and V. Berestetskii, Quantum Electrodynamics, New York, (1965).

[4] D. Sivers, Phys. Rev. D **41**, 83 (1990).

[5] D. Sivers, Phys. Rev. D **43**, 261 (1991).

[6] V. N. Baier, E. A. Kuraev, V. S. Fadin, and V. A. Khoze, Phys. Rept. **78**, 293 (1981).

[7] V. M. Budnev, I. F. Ginzburg, G. V. Meledin, and V. G. Serbo, Phys. Rept. **15**, 181 (1975).

[8] S. Weinberg, Phys. Rev. **150**, 1313 (1966).

[9] Dr. A.P. Nagaytsev, Private communication.